

Lecture Notes, January 20, 2012

Convexity

A set of points S in R^N is said to be convex if the line segment between any two points of the set is completely included in the set.

S is convex if $x, y \in S$, implies $\{z \mid z = \alpha x + (1 - \alpha)y, 0 \leq \alpha \leq 1\} \subseteq S$.

S is said to be strictly convex if $x, y \in S, x \neq y, 0 < \alpha < 1$, implies $\alpha x + (1 - \alpha)y \in \text{interior } S$.

The notion of convexity is that a set is convex if it is connected, has no holes on the inside and no indentations on the boundary. A set is strictly convex if it is convex and has a continuous strict curvature (no flat segments) on the boundary.

Economically, this notion corresponds to "diminishing marginal utility"
"diminishing marginal rate of substitution" "diminishing marginal product" .

Properties of Convex Sets

Let C_1, C_2 be convex subsets of R^N . Then:

$C_1 \cap C_2$ is convex,

$C_1 + C_2$ is convex,

$\overline{C_1}$ is convex

The unit simplex in \mathbf{R}^N , is

$$P = \left\{ p \mid p \in \mathbf{R}^N, p_i \geq 0, i = 1, \dots, N, \sum_{i=1}^N p_i = 1 \right\}. \quad (5.1)$$

The unit simplex is a (generalized) triangle in N -space.

Note that P is compact (closed and bounded) and convex.

Theorem 5.1 (Brouwer Fixed-Point Theorem) Let $f(\cdot)$ be a continuous function, $f : P \rightarrow P$. Then there is $x^* \in P$ so that $f(x^*) = x^*$.

The four properties assumed in the Brouwer Fixed Point Theorem — continuity of f , closedness, boundedness, and convexity of P — are all essential to the theorem. Omit any one of them and the theorem fails.

The result can be generalized from P to any compact convex set.